

KCL-TH-92-2

July 1992

On the quantum KP hierarchy and its relation to the non-linear Schrödinger equation

M.D. Freeman and P. West

King's College, Department of Mathematics, Strand, London WC2R 2LS, GB

Abstract

We establish a relation between the classical non-linear Schrödinger equation and the KP hierarchy, and we extend this relation to the quantum case by defining a quantum KP hierarchy. We also present evidence that an integrable hierarchy of equations is obtained by quantizing the first Hamiltonian structure of the KdV equation. The connection between infinite-dimensional algebras and integrable models is discussed.

1 Introduction

Integrable dynamical systems with an infinite number of degrees of freedom have been much studied for many years. For such dynamical systems, integrability is taken to mean the existence of an infinite number of conserved quantities in involution, i.e. for which the Poisson bracket between any pair is zero. Examples of such systems include the KdV equation,

$$\frac{\partial u}{\partial t} = \frac{1}{8}u''' - \frac{3}{8}uu', \quad (1.1)$$

the Liouville equation,

$$\partial\bar{\partial}\phi - e^{-\phi} = 0, \quad (1.2)$$

and the sinh-Gordon equation

$$\partial\bar{\partial}\phi + 2\sinh\phi = 0. \quad (1.3)$$

These systems are in fact related to the Lie algebra $sl(2)$. Corresponding to every Lie algebra there is an integrable generalization of each of them [1, 2], these being the generalized KdV equations, Toda equations and affine Toda equations respectively.

All of these equations can be written in Hamiltonian form. In each case the Hamiltonian generating the time evolution has the property that it commutes with all of the conserved charges with respect to the Poisson bracket. Each equation is therefore a member of an infinite hierarchy of equations, obtained by using the conserved charges as Hamiltonians to define commuting time evolutions for infinitely many times.

In another development, which at first sight has no connection with integrability, conformal field theories have been intensively studied recently. A theory that is Poincaré invariant possesses a conserved energy-momentum tensor, and if such a theory is in addition conformally invariant then the energy-momentum tensor is traceless. In two dimensions this implies that, in light-cone coordinates $(z, \bar{z})$, the component $T \equiv T_{zz}$ is independent of $\bar{z}$,

$$\bar{\partial}T = 0, \quad (1.4)$$

and so any polynomial in T is conserved. Thus the integrals of such polynomials are conserved charges for the theory.

It is a remarkable fact that all the integrable systems referred to above can be linked to the infinite-dimensional conformal algebra and its extensions. Of course any conformally invariant theory in two dimensions automatically possesses an infinite number of conserved quantities, as argued above, but in order to have integrability we require a set of conserved quantities in involution. Hence it is necessary to select from the set of all polynomials in T a subset for which the Poisson bracket between any pair of charges is zero. It turns out that the charges arrived at in this way are essentially the conserved charges of the sinh-Gordon and KdV equations, even though these systems are not conformally invariant. A similar result holds for the generalizations of these systems corresponding to arbitrary Lie algebras. For each Lie algebra there is an extension of the conformal algebra, referred to generically as a W -algebra, and choosing a commuting subset of conserved quantities in the W -algebra corresponding to the Lie algebra $\mathfrak{g}$ gives the conserved charges for the integrable systems corresponding to $\mathfrak{g}$.

Let us spell out in more detail how this procedure works in the simplest case. The first step is to write the KdV equation in Hamiltonian form. It is well known that there are two possible Poisson brackets for the field $u(x)$. With respect to the so-called second Poisson bracket structure

$$\{u(x), u(y)\} = \left(u(x)\partial_x + \partial_x u(x) - \partial_x^3 \right) \delta(x - y) \quad (1.5)$$

the time evolution is given by the Hamiltonian

$$H_2 = \frac{1}{16} \int u^2, \quad (1.6)$$

while in terms of the first Poisson bracket structure

$$\{u(x), u(y)\} = \partial_x \delta(x - y) \quad (1.7)$$

the time evolution is given by

$$H_3 = -\frac{1}{16} \int u^3 + u'^2. \quad (1.8)$$

It is through the second Poisson bracket that the connection with the conformal algebra is made. It was recognized in [3] that the second Poisson bracket for the Fourier modes of u is precisely the two-dimensional conformal algebra, so that we can identify u with the energy-momentum tensor T . The conserved quantities of the KdV equation are then integrals of polynomials in T and its derivatives.

If now we take a realization of the second Poisson bracket structure of the KdV equation in terms of some field Φ , so that u is expressed in terms of Φ and the Poisson bracket for u follows from that of Φ , we can obtain an integrable hierarchy of evolution equations for Φ . This hierarchy is obtained simply by taking the conserved charges for the KdV equation, expressing them in terms of Φ , and then using these as Hamiltonians to generate time evolutions for Φ . Such a hierarchy is obtained for every possible realization of the Virasoro algebra.

An example of this is the realization of the second Poisson bracket structure of the KdV equation given by the Miura transformation. The field $u(x)$ is expressed in terms of a field $v(x)$ satisfying

$$\{v(x), v(y)\} = \partial_x \delta(x - y) \quad (1.9)$$

through the transformation

$$u = \frac{1}{2}v^2 + v'. \quad (1.10)$$

The hierarchy of evolution equations obtained in this way for $v(x)$ is precisely the mKdV hierarchy. We can take this example one stage further by taking v to be the derivative of a scalar field ϕ . In this case the Poisson bracket of u with $\int dx \exp(-\phi(x))$ is easily seen to be zero. In fact the conserved quantities of the KdV equation, when expressed in terms of v through the Miura transformation, are even functions of v , and so these charges must also commute with $\int dx \exp(\phi(x))$ and hence with $\int dx \cosh \phi(x)$. But $\int dx \cosh \phi(x)$ is the Hamiltonian for the sinh-Gordon equation, and so the conserved charges for the KdV equation are also conserved for the sinh-Gordon equation. Hence the KdV, mKdV and sinh-Gordon equations have essentially the same conserved quantities because their fields provide different realizations of the classical Virasoro algebra.

There are analogues of this picture for other integrable systems. In particular let us consider the N 'th KdV hierarchy, for variables $u_i, i = 1, \dots, N$, corresponding to the Lie algebra $sl(N + 1)$. The equations of this hierarchy can be written in Hamiltonian form with Poisson brackets that are a classical limit of the $\mathbf{W}_{(N+1)}$ algebra of Fateev and Lukyanov [4]. There is a realization of the u_i in terms of fields ϕ_j with Poisson brackets $\{\phi'_i(x), \phi'_j(y)\} = \delta'(x - y)$ via the generalized Miura transformation

$$\prod_i (\partial + h_i \cdot \phi) = \sum u_s \partial^{N-s}, \quad (1.11)$$

where the h_i are the weights of the fundamental representation of $sl(N+1)$. The evolution equations for the ϕ_i generated by the Hamiltonians of the N 'th KdV hierarchy are those of the N 'th mKdV hierarchy, and these equations have the same conserved quantities as the affine Toda field theory based on $sl(N+1)$.

The above considerations have all been classical, and it is natural to consider whether the same picture holds after quantization. The quantum KdV, quantum mKdV and quantum sinh-Gordon equations were considered in [5], and the first few commuting conserved quantities were constructed. It was indeed found that for these quantities the connection spelt out above held also at the quantum level. Thus the charges for the quantum KdV equation, when expressed in terms of a scalar field through the Feigin-Fuchs construction, were precisely the charges found for the quantum mKdV and sinh-Gordon equations. The first few conserved charges for these systems have also been found from the perspective of perturbations of conformal field theories [6, 7, 8]. The existence of conserved commuting charges at all expected levels has argued in a free-field realization in [9], but the methods that are used classically to construct the conserved charges and establish their properties have not so far been carried over to the quantum theories. The difficulties in doing this can be traced to the necessity of normal-ordering the quantum conserved currents and their products.

One approach to establishing the integrability of quantum theories would be to attempt to exploit the remarkable connection spelt out above at the classical level between infinite-dimensional algebras and integrability. Indeed one might conjecture that this pattern is the general case, and that every integrable system is associated with the Virasoro algebra, or an extension of it, in this way. Then different realizations of a given algebra will lead to different integrable systems, but all such systems originating from a given algebra will have the same conserved quantities.

Clearly, given an infinite set of commuting quantities, we can define an integrable system by taking them as Hamiltonians. In this sense one can regard the infinite set of commuting quantities as the primary object. From this point of view the conjecture is that any such set of quantities is associated with the Virasoro algebra or a W -algebra. One can also go further and conjecture that the commuting quantities can be seen as a Cartan subalgebra of an even larger algebra. Some evidence for this point of view was given in [10], where

a search for commuting charges constructed from a $U(1)$ Kac-Moody current was carried out. An infinite number of sets of such commuting charges was seen to exist, but it was found that, although the calculation was carried out entirely within the framework of the $U(1)$ Kac-Moody algebra, each set of commuting charges could be associated with an exceptional W -algebra.

The motivation for this paper was to consider other integrable systems and examine the extent to which they conformed with the above conjecture. To this end we examined the non-linear Schrödinger equation. We will show that there is a relation between this equation and the KP hierarchy which is analagous to that which exists between the mKdV and KdV hierarchies. The essence of the connection between the non-linear Schrödinger equation and the KP hierarchy is that the first Poisson bracket structure [11] of the KP hierarchy is isomorphic [12, 13] to the algebra $W_{1+\infty}$, which is a linear extension of the Virasoro algebra containing a single spin- i quasiprimary field for each spin $i \geq 1$ [14], and this algebra has a realization in terms of a complex scalar field ψ, ψ^* .

The quantum KP hierarchy is then defined using the quantum $W_{1+\infty}$ algebra, and it is shown to possess conserved quantities at least at the first few levels. The connection between the classical non-linear Schrödinger equation and the KP hierarchy spelt out above is shown to extend to the quantum analogues.

2 The relation between the KP hierarchy and the non-linear Schrödinger equation

The non-linear Schrödinger equation is given by

$$\frac{\partial \psi}{\partial t} + \psi'' + 2\kappa \psi^* \psi^2 = 0, \quad (2.1)$$

where ψ, ψ^* is a bosonic complex scalar field and κ is a coupling constant. This equation can be written in Hamiltonian form $\dot{\psi} = \{\psi, H\}$ provided we adopt the Poisson brackets

$$\{\psi^*(x), \psi(y)\} = \delta(x - y), \quad \{\psi(x), \psi(y)\} = \{\psi^*(x), \psi^*(y)\} = 0 \quad (2.2)$$

and take $H = \int (\psi^* \psi'' + \kappa \psi^* \psi^2)$.

It is useful to note that it is possible to assign weights to ψ and ψ^* in a manner consistent with the Poisson bracket. We take ∂ and $\delta(x - y)$ to have weight 1, and then eqn (2.2) implies that the product $\psi^*\psi$ must have weight 1. If we now take κ to have weight 1, the Hamiltonian H will be homogeneous of weight 2.

The non-linear Schrödinger equation possesses an infinite set of conserved quantities Q_n , given by the formula

$$Q_n = \frac{1}{\kappa} \int dx \psi(x) Y_n(\psi, \psi^*), \quad (2.3)$$

where

$$Y_1 = \kappa \psi^*, \quad Y_{n+1} = Y'_n + \psi \sum_{k=1}^{n-1} Y_k Y_{n-k} \quad \text{for } n \geq 1. \quad (2.4)$$

The first few such charges are

$$\begin{aligned} Q_1 &= \int dx \psi^* \psi \\ Q_2 &= - \int dx \psi^* \psi' \\ Q_3 &= \int dx (\psi^* \psi'' + \kappa \psi^{*2} \psi^2); \end{aligned} \quad (2.5)$$

it can be checked that these charges commute and so can be used to generate commuting evolutions, giving a hierarchy of equations.

The KP hierarchy consists of an infinite set of differential equations which has, at first sight, no relation to the non-linear Schrödinger equation. This set of equations is defined in terms of a pseudodifferential operator Q of the form

$$Q = D + q_0 D^{-1} + q_1 D^{-2} + \dots \quad (2.6)$$

Here D denotes $\partial/\partial z$ and satisfies the following generalization of the Leibniz rule:

$$D^n f = \sum_{r=0}^{\infty} \binom{n}{r} f^{(r)} D^{n-r}, \quad n \in \mathbb{Z}. \quad (2.7)$$

An infinite set of commuting time evolutions for the fields q_i , $i = 0, 1, 2, \dots$, is given by

$$\frac{\partial Q}{\partial t_p} = [(Q^p)_+, Q], \quad p = 0, 1, 2, \dots \quad (2.8)$$

where $(Q^p)_+$ is the part of Q^p involving no negative powers of D . These evolution equations can be written in Hamiltonian form. There are in fact a number of ways to do this,

but for our purposes it is the so-called first Hamiltonian structure that is needed. The Poisson bracket in this case is obtained from the method of coadjoint orbits applied to the Lie algebra of differential operators. For convenience we summarize the essential elements of this method as applied to a general Lie algebra $\mathfrak{g}$. The idea is to define Poisson brackets between functions on the dual space $\mathfrak{g}^*$ of $\mathfrak{g}$. Every element X of the Lie algebra $\mathfrak{g}$ defines a natural linear function on $\mathfrak{g}^*$ according to the formula

$$X(\omega) \equiv \omega(X), \quad \forall \quad \omega \in \mathfrak{g}^*. \quad (2.9)$$

Then, given $X, Y \in \mathfrak{g}$, thought of as functions on $\mathfrak{g}^*$, we define their Poisson bracket to be the function on $\mathfrak{g}^*$ given by

$$\{X, Y\}(\omega) = \omega([X, Y]). \quad (2.10)$$

Thus, on the left-hand side of this equation X and Y are considered as functions on $\mathfrak{g}^*$, while on the right-hand side they are thought of as elements of the Lie algebra $\mathfrak{g}$.

In the case at hand $\mathfrak{g}$ is taken to be the Lie algebra of differential operators, and the dual $\mathfrak{g}^*$ can be identified with the space of pseudodifferential operators of the form (2.6). Given a differential operator X , which we write for convenience in the form $X = x_0(z) + D x_1(z) + \dots + D^n x_n(z)$, where the derivative $D \equiv \frac{\partial}{\partial z}$ acts to the right, and a pseudodifferential operator Q as in eqn (2.6), we take

$$\begin{aligned} Q(X) &\equiv \text{tr}(QX) \\ &\equiv \int dz \{ \text{coefficient of } D^{-1} \text{ in } QX \} \\ &= \int dz \sum_{i=0}^n q_i(z) x_i(z). \end{aligned} \quad (2.11)$$

Taking $X(z) = D^i \delta(z - x)$ and $Y(z) = D^j \delta(z - y)$, we obtain

$$\begin{aligned} \{q_i(x), q_j(y)\} &\equiv \{X, Y\}(Q) \\ &= Q([X, Y]) \\ &= \sum_{r=0}^j \binom{j}{r} \partial_x^r (q_{i+j-r}(x) \delta(x - y)) - \sum_{r=0}^i \binom{i}{r} (-1)^r q_{i+j-r} \partial_x^r \delta(x - y). \end{aligned} \quad (2.12)$$

In evaluating the commutator $[X, Y]$ we made use of the relation

$$f D^m = \sum_{r=0}^m (-1)^{m-r} \binom{m}{r} D^r f^{(m-r)}. \quad (2.13)$$

The algebra (2.13) is the first Poisson bracket structure for the KP hierarchy [11]. This algebra is in fact precisely $\mathbf{W}_{1+\infty}$ with zero central charge [12, 13]; the q_i are related to the usual basis for $\mathbf{W}_{1+\infty}$, denoted by V^i [14, 15], by a linear transformation. It can be shown that the time evolutions (2.8) can be written as

$$\frac{\partial q_i}{\partial t_p} = \{q_i, H_{p+1}\}, \quad p = 0, 1, 2, \dots, \quad (2.14)$$

where

$$H_p = \frac{1}{p} \text{tr}(Q^p) \quad (2.15)$$

Just as for the non-linear Schrödinger equation, there is an assignment of weights to the q_i which is consistent with the Poisson bracket structure (2.13). If we take ∂ and $\delta(x-y)$ to have weight 1, then q_i must have weight $i+1$. With this choice, the operator Q in eqn (2.6) does not have a definite weight, but we can correct this by working instead with

$$\tilde{Q} \equiv \kappa^{-1} D + q_0 D^{-1} + q_1 D^{-2} + \dots, \quad (2.16)$$

where, as before, κ is a constant of weight 1. $\tilde{Q}$ then has weight zero. Using $\tilde{Q}$ instead of Q does not change the Poisson brackets between the q_i , but the Hamiltonians

$$\tilde{H}_p = \frac{1}{p} \text{tr}(\tilde{Q}^p) \quad (2.17)$$

now involve various powers of κ . The first few $\tilde{H}_p$ are given explicitly by

$$\begin{aligned} \tilde{H}_1 &= \int dx q_0 \\ \tilde{H}_2 &= \kappa^{-1} \int dx q_1 \\ \tilde{H}_3 &= \kappa^{-2} \int dx (q_2 + \kappa q_0^2) \\ \tilde{H}_4 &= \kappa^{-4} \int dx (q_3 + 3\kappa q_0 q_1). \end{aligned} \quad (2.18)$$

The $\tilde{H}_p$ can be viewed as a set of Poisson commuting quantities within the enveloping algebra of $\mathbf{W}_{1+\infty}$.

Having given the KP and the non-linear Schrödinger equations, we are now in a position to explain the relation between them by exploiting the fact that $\mathbf{W}_{1+\infty}$ has a realization in terms of a single complex boson. Let us take a general approach first. We consider a

realization of $W_{1+\infty}$ in terms of some set of fields ϕ_i , so that the q_i can be expressed in terms of the ϕ_i and the Poisson brackets for the q_i follow from those of the ϕ_i . If such a realization of $W_{1+\infty}$ has zero central charge, we can clearly construct an infinite set of mutually commuting quantities by expressing the H_n 's in terms of the ϕ_i . We can then define time evolutions for the ϕ_i by

$$\frac{\partial \phi_i}{\partial t_p} = \{\phi_i, H_{p+1}\}, \quad p = 0, 1, 2, \dots, \quad (2.19)$$

and it follows from the Leibniz property of the Poisson bracket that this implies the KP evolutions for the q_i .

There is a centreless realization of $W_{1+\infty}$ in terms of a complex boson with Poisson brackets given by

$$\{\psi^*(x), \psi(y)\} = \delta(x - y), \quad \{\psi(x), \psi(y)\} = \{\psi^*(x), \psi^*(y)\} = 0. \quad (2.20)$$

This is most easily seen by checking that the correct Poisson bracket (2.13) between the q_i is obtained by taking

$$q_i = (-1)^i \psi^* \psi^{(i)}. \quad (2.21)$$

The first few Hamiltonians of the KP hierarchy, when expressed in terms of ψ and ψ^* , take the forms

$$\begin{aligned} \tilde{H}_1(\psi^*, \psi) &= \int dx \psi^* \psi \\ \tilde{H}_2(\psi^*, \psi) &= -\kappa^{-1} \int dx \psi^* \psi' \\ \tilde{H}_3(\psi^*, \psi) &= \kappa^{-2} \int dx (\psi * \psi'' + \kappa \psi^{*2} \psi^2) \\ \tilde{H}_4(\psi^*, \psi) &= -\kappa^{-4} \int dx (\psi^* \psi''' + 3\kappa \psi^{*2} \psi \psi'). \end{aligned} \quad (2.22)$$

We recognize these immediately as the first few conserved charges of the non-linear Schrödinger equation, up to overall factors. It follows from the commutativity of the KP Hamiltonians that the $\tilde{H}_n(\psi^*, \psi)$ will also commute. However, since the charges of the non-linear Schrödinger equation are uniquely determined by the requirement that they commute with $\tilde{H}_3$, it follows that the charges obtained from the KP system must coincide with those of the non-linear Schrödinger equation at all levels.

3 A quantum KP hierarchy

Given that the classical non-linear Schrödinger equation is related to the KP hierarchy and that the quantum non-linear Schrödinger equation is known to be integrable, one might expect there to exist an integrable quantum KP hierarchy. One way to construct such a hierarchy would be to take the first Poisson bracket structure for the KP hierarchy, which is isomorphic to $\mathbf{W}_{1+\infty}$, and then to replace Poisson brackets by commutators. The existence of an integrable quantum KP hierarchy would then follow if we could find an infinite set of commuting Hamiltonians in the enveloping algebra of $\mathbf{W}_{1+\infty}$.

In this section we construct some quantum analogues of the Hamiltonians $\tilde{H}_p$ from the generators V^i of $\mathbf{W}_{1+\infty}$. We use the notation of [14], where V^i is a quasiprimary field of weight $i + 2$. The first step is to obtain explicit expressions for the operator product expansions for the V^i , which are given in terms of generalized hypergeometric functions in [15]. We used Mathematica [16] to calculate these hypergeometric functions; the first few operator product expansions are given by

$$\begin{aligned}
V^{-1}(z)V^{-1}(w) &= \frac{c}{(z-w)^2} + \dots \\
V^{-1}(z)V^0(w) &= \frac{V^{-1}(w)}{(z-w)^2} + \dots \\
V^0(z)V^0(w) &= \frac{c/2}{(z-w)^4} + \frac{2V^0(w)}{(z-w)^2} + \frac{V^{0'}(w)}{z-w} + \dots \\
V^{-1}(z)V^1(w) &= \frac{2V^0(w)}{(z-w)^2} + \dots \\
V^0(z)V^1(w) &= \frac{V^{-1}(w)}{(z-w)^4} + \frac{3V^1(w)}{(z-w)^2} + \frac{V^{1'}(w)}{z-w} + \dots
\end{aligned} \tag{3.1}$$

We then looked for commuting charges that could be expressed as integrals of sums of products of the fields in $\mathbf{W}_{1+\infty}$. The commutator of such charges is found in the usual way from the operator product expansions; the integrals of fields A and B commute if the single pole term in the operator product expansion of A with B is a derivative. We used the package OPEdefs of Thielemans [17] to calculate the necessary OPE's. The first two charges, namely

$$Q_1 = \int dz V^{-1} \tag{3.2}$$

and

$$Q_2 = \int dz V^0 \quad (3.3)$$

are trivial, in the sense that the first of these commutes with all $V^i(z)$ while the commutator of Q_2 with $V^i(z)$ is just $V^{i'}(z)$. For the first non-trivial charge we take

$$Q_3 \equiv \int dz \{V^1 + \kappa (V^{-1})^2\}; \quad (3.4)$$

this is analogous to the classical Hamiltonian H_3 of the KP hierarchy in that it involves terms of weights 2 and 3. In this and subsequent expressions involving products of operators, we use the normal ordering standard in conformal field theory, namely we take

$$(AB)(z) = \oint_z dw \frac{A(w)B(z)}{w - z}. \quad (3.5)$$

It is our belief that there are infinitely many further charges that commute with the charge Q_3 , and that furthermore these charges are unique. Using a computer we have found three such charges, and we have verified that all the charges we have obtained commute amongst themselves. We conjecture that there exists an infinite set of such charges, one for each spin. We give here Q_4 and Q_5 :

$$\begin{aligned} Q_4 &= \int dz \{V^2 + 3\kappa (V^{-1}V^0) + 3c\kappa^2/2 (V^{-1})^2\} \\ Q_5 &= \int dz \{V^3 + \kappa (4(V^{-1}V^1) + 2(V^0)^2 - 40/3 (V^{-1})^2) \\ &\quad + \kappa^2 (2(V^{-1})^3 + 4c(V^{-1}V^0)) + 2c^2\kappa^3 (V^{-1})^2\}. \end{aligned} \quad (3.6)$$

4 The quantum non-linear Schrödinger equation

The integrability of the quantum non-linear Schrödinger equation has been extensively studied using the quantum inverse scattering method and also from the point of view of the existence of an infinite number of commuting conserved quantities. The relation between these different approaches was discussed in [18]. In order to quantize the non-linear Schrödinger equation we adopt the following OPE for the bosonic operators ψ and ψ^* :

$$\psi^*(z)\psi(w) = (z - w)^{-1} + \dots \quad (4.1)$$

The time evolution is generated by

$$H = \oint dz (: \psi^* \psi'' : + : \kappa \psi^{*2} \psi^2 :), \quad (4.2)$$

in the sense that

$$\frac{\partial \psi}{\partial \bar{z}} = [\psi, H], \quad \frac{\partial \psi^*}{\partial \bar{z}} = [\psi^*, H]. \quad (4.3)$$

We can then search for normal-ordered polynomials in ψ, ψ^* and their derivatives whose integrals commute with H ; the first few conserved currents are as follows:

$$\psi^* \psi, \quad \psi^* \psi', \quad \psi^* \psi'' + \kappa \psi^{*2} \psi^2, \quad \psi^* \psi''' + 6\kappa \psi^* \psi'' + 3\kappa \psi^{*2} \psi \psi'. \quad (4.4)$$

In section 2 we showed that the non-linear Schrödinger equation and the KP hierarchy were related at the classical level, and we now extend that analysis to the quantum case. Just as the Poisson algebra $\mathbf{W}_{1+\infty}$ has a realization in terms of classical scalar fields ψ and ψ^* , the quantum algebra $\mathbf{W}_{1+\infty}$ has a realization in terms of operators ψ and ψ^* with OPE given by eqn (4.1). The quasiprimary fields V^i of $\mathbf{W}_{1+\infty}$ have the form

$$V^i = \binom{2n+2}{n+1}^{-1} \sum_{r=0}^{i+1} (-1)^r \binom{i+1}{r}^2 \psi^{(r)} \psi^{*(i+1-r)}. \quad (4.5)$$

On substituting these expressions into the quantum KP charges Q_i of the previous section we recover the charges of the quantum non-linear Schrödinger equation given above, up to the freedom to add lower spin charges. Some care must be taken in transforming from the normal ordering in terms of the V^i to the free field normal ordering used for the non-linear Schrödinger charges. Since the KP charges commute it follows that the non-linear Schrödinger charges will also commute, and because the non-linear Schrödinger and KP charges are uniquely determined by requiring that they commute with H_3 we conclude that we obtain all the non-linear Schrödinger charges in this way.

5 Quantizing the first Hamiltonian structure for the KdV equation

We have seen that there is strong evidence that quantizing the first Hamiltonian structure of the KP hierarchy leads to a quantum integrable system. This leads one to suspect that

it may be possible to define an alternative integrable quantum KdV system by quantizing the first Poisson bracket structure $\{u(x), u(y)\} = \delta'(x - y)$ for this equation. We demand that u be the current of a $U(1)$ Kac-Moody algebra and so satisfy the OPE

$$u(z)u(w) = (z - w)^{-2} + \dots \quad (5.1)$$

We consider a time evolution generated by $H = \oint (u'^2 + \kappa u^3)$, where the coupling constant κ is included so that H has definite weight, and look for charges commuting with this operator. We have found three non-trivial charges commuting with H . These charges also commute with each other. The first two are

$$\oint dz \left\{ u''^2 + 5\kappa uu'^2 + 5\kappa^2/4 (u^4 - u'^2) \right\} \quad (5.2)$$

$$(5.3)$$

the third is a polynomial of degree 7 in κ , with first term u''''^2 . The charges found in this way are not the same as those obtained by quantizing the second Poisson bracket structure of the KdV equation. It has already been noted [19] that the first non-trivial equations of motion obtained from the first and second Poisson bracket structures are not equivalent in the quantum case. It is remarkable, however, that each appears to lead to an integrable quantum hierarchy.

6 Conclusions

We have shown that the classical non-linear Schrödinger equation is obtained as a particular realization of the KP hierarchy, and by quantizing the first Hamiltonian structure of the KP hierarchy we have shown that the quantum non-linear Schrödinger equation can be obtained in a similar way. It would be of great interest to quantize the second Hamiltonian structure of the KP hierarchy. The second Poisson bracket structure is known [20], but it is nonlinear, and its quantum analogue has not been found. Given this latter structure, however, it would be straightforward to quantize the KP hierarchy by taking commuting charges as Hamiltonians. It would be interesting to study the quantum Hamiltonian reduction of the quantum KP hierarchy to obtain the quantum KdV equation; such a step would explain the structure of the charges and possibly the integrability of the latter

equation. It would also be interesting to investigate further the relationship between the two quantum integrable systems obtained by quantizing the two Hamiltonian structures for the KdV equation.

Acknowledgement: We are grateful to Klaus Hornfeck for many useful discussions. M.D. Freeman is grateful to the UK Science and Engineering Research Council for financial support.

Note added: After this work was completed, Dr J. Gibbons informed us of a connection between the non-linear Schrödinger equation and the KdV equation. We would like to thank him for this information.

References

- [1] V.G. Drinfeld and V.V. Sokolov, J. Sov. Math. **30** (1984) 1975.
- [2] G. Wilson, Ergod. Th. and Dynam. Sys. **1** (1981) 361.
- [3] J.-L. Gervais, Phys. Lett. B160 (1985) 277; Phys. Lett. B160 (1985) 279.
- [4] V.A. Fateev and S.L. Lukyanov, Int. J. Mod. Phys. A3 (1988) 507; “Additional symmetries and exactly-soluble models in two-dimensional conformal field theory,” Moscow preprint 1988.
- [5] R. Sasaki and I. Yamanaka, Adv. Studies in Pure Math. **16** (1988) 271.
- [6] A.B. Zamolodchikov, “Integrable field theory from conformal field theory,” Proceedings of the Taniguchi Symposium, Kyoto, (1988); Int. J. Mod. Phys. A3 (1988) 743; Int. J. Mod. Phys. A4 (1989) 4235.
- [7] T. Eguchi and S-K. Yang, Phys. Lett. B224 (1989) 373.
- [8] T. Hollowood and P. Mansfield, Phys. Lett. B226 (1989) 73.
- [9] B. Feigin and E. Frenkel, Phys. Lett. B276 (1992) 79.
- [10] M.D. Freeman, K. Hornfeck and P. West, “Commuting quantities and exceptional W-algebras,” to be published in Int. J. Mod. Phys. A

- [11] Y. Watanabe, Lett. Math. Phys. 7 (1983) 99.
- [12] K. Yamagishi, Phys. Lett. B259 (1991) 436.
- [13] F. Yu and Y-S. Wu, Phys. Lett. B263 (1991) 220.
- [14] C.N. Pope, L.J. Romans and X. Shen, Phys. Lett. B242 (1990) 401.
- [15] C.N. Pope, “Lectures on W algebras and W gravity,” preprint CTP TAMU-103/91.
- [16] S. Wolfram, “Mathematica: a system for doing mathematics by computer,” Addison-Wesley, 1991.
- [17] K. Thielemans, Int. J. Mod. Phys. **C2** (1991) 787.
- [18] M. Omote, M. Sakagami, R. Sasaki and I. Yamanaka, Phys. Rev. D35 (1987) 2423.
- [19] B.A. Kupershmidt and P. Mathieu, Phys. Lett. B227 (1989) 245.
- [20] L.A. Dickey, Ann. N.Y. Acad. Sci. 491 (1987) 131; J.M. Figueroa-O’Farrill, J. Mas and E. Ramos, Phys. Lett. B266 (1991) 298; F. Yu and Y-S. Wu, Nucl. Phys. B373 (1992) 713; A. Das, W-J Huang and S. Panda, Phys. Lett. B271 (1991) 109.